

GROUP	<b>12</b>	FUNGICIDE
-------	-----------	-----------

## MEDALLION® Fungicide

**FUNGICIDE**

**SUSPENSION**

**COMMERCIAL**

For control of diseases on turf. For control and suppression of diseases in outdoor and greenhouse ornamentals, and greenhouse vegetables (cucumber, peppers and tomatoes).

**ACTIVE INGREDIENT:**

Fludioxonil ..... 125 g/L

Contains 1,2-benzisothiazolin-3-one at 0.037% and 2-bromo-2-nitropropane-1,3-diol at 0.036% OR 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00091%, 2-methyl-4-isothiazolin-3-one at 0.00030% and 2-bromo-2-nitropropane-1,3-diol at 0.036% as preservatives.

**READ THE LABEL AND PAMPHLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO.: **31528**  
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **0.5 L - Bulk**

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario N1G 4Z3  
Telephone: 1-877-964-3682

Label

## 1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

## 2.0 FIRST AID

**IF POISONING IS SUSPECTED, IMMEDIATELY** contact a physician or a poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## 3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote known. Treat symptomatically.

## 4.0 PRECAUTIONS

**KEEP OUT OF REACH OF CHILDREN.**

**DO NOT** enter or allow entry into treated turf areas until MEDALLION® Fungicide is dry. For all other crops, **DO NOT** enter or allow worker entry into treated areas for 12 hours following application.

## 5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls or a long sleeved shirt and long pants, chemical resistant gloves, and goggles when mixing, loading or during equipment clean up or repair. Wear coveralls or a long sleeved shirt and long pants during application.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean

clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

## 6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Fludioxonil is persistent and may carryover. It is recommended that any products containing fludioxonil not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

## 7.0 STORAGE

Store in original container in a cool, dry place. **DO NOT** use or store near heat or open flame. **DO NOT** store this product under wet conditions. Keep container closed when not in use. To prevent contamination, store this product away from food or feed.

## 8.0 DISPOSAL

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or provincial regulatory agency.

### For Recyclable Containers

**DO NOT** reuse this container for any other purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank or dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**For Refillable Containers**

**DO NOT** reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

**For Returnable Containers**

**DO NOT** reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (1-800-FASTMED)***

MEDALLION® is a trademark of a Syngenta Group Company.

GROUP	<b>12</b>	FUNGICIDE
-------	-----------	-----------

## MEDALLION® Fungicide

**FUNGICIDE**

**SUSPENSION**

**COMMERCIAL**

For control of diseases on turf. For control and suppression of diseases in outdoor and greenhouse ornamentals, and greenhouse vegetables (cucumber, peppers and tomatoes).

**ACTIVE INGREDIENT:**

Fludioxonil ..... 125 g/L

Contains 1,2-benzisothiazolin-3-one at 0.037% and 2-bromo-2-nitropropane-1,3-diol at 0.036% OR 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00091%, 2-methyl-4-isothiazolin-3-one at 0.00030% and 2-bromo-2-nitropropane-1,3-diol at 0.036% as preservatives.

**READ THE LABEL AND PAMPHLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

REGISTRATION NO.: **31528**  
PEST CONTROL PRODUCTS ACT

**Syngenta Canada Inc.**  
140 Research Lane, Research Park  
Guelph, Ontario N1G 4Z3  
Telephone: 1-877-964-3682

Pamphlet

## TABLE OF CONTENTS

	<u>Section Number</u>
NOTICE TO USER .....	1.0
FIRST AID .....	2.0
TOXICOLOGICAL INFORMATION .....	3.0
PRECAUTIONS .....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE) .....	5.0
ENVIRONMENTAL PRECAUTIONS .....	6.0
STORAGE .....	7.0
DISPOSAL .....	8.0
PRODUCT INFORMATION .....	9.0
DIRECTIONS FOR USE .....	10.0
General Information .....	10.1
Application Instructions .....	10.2
Mixing Instructions .....	10.3
MEDALLION Fungicide and Tank Mixtures .....	10.4
Buffer Zones .....	10.5
CROP USE DIRECTIONS .....	11.0
Turf: Golf Courses .....	11.1
MINOR USES .....	12.0
Ornamentals (Outdoor and Greenhouse) .....	12.1
Greenhouse Vegetables .....	12.2
RESISTANCE MANAGEMENT RECOMMENDATIONS .....	13.0

## 1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

## 2.0 FIRST AID

**IF POISONING IS SUSPECTED, IMMEDIATELY** contact a physician or a poison control centre. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED**, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING**, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## 3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote known. Treat symptomatically.

## 4.0 PRECAUTIONS

**KEEP OUT OF REACH OF CHILDREN.**

**DO NOT** enter or allow entry into treated turf areas until MEDALLION® Fungicide is dry. For all other crops, **DO NOT** enter or allow worker entry into treated areas for 12 hours following application.

## 5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls or a long sleeved shirt and long pants, chemical resistant gloves, and goggles when mixing, loading or during equipment clean up or repair. Wear coveralls or a long sleeved shirt and long pants during application.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean

clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

## 6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Fludioxonil is persistent and may carryover. It is recommended that any products containing fludioxonil not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

## 7.0 STORAGE

Store in original container in a cool, dry place. **DO NOT** use or store near heat or open flame. **DO NOT** store this product under wet conditions. Keep container closed when not in use. To prevent contamination, store this product away from food or feed.

## 8.0 DISPOSAL

For information on disposal of unused, unwanted product, or in the case of a spill or spill clean-up, contact the manufacturer or provincial regulatory agency.

### For Recyclable Containers

**DO NOT** reuse this container for any other purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank or dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **For Refillable Containers**

**DO NOT** reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

### **For Returnable Containers**

**DO NOT** reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,  
CALL 1-800-327-8633 (1-800-FASTMED)***

## **9.0 PRODUCT INFORMATION**

MEDALLION Fungicide is a broad spectrum contact and protective fungicide for control of listed diseases in turf and control and suppression of diseases in outdoor and greenhouse ornamentals and suppression of diseases in greenhouse vegetables. It is a suspension concentrate formulation containing 125 g/L fludioxonil. Fludioxonil is a long lasting contact fungicide belonging to the phenyl pyrrole chemistry group that provides broad-spectrum activity against a wide range of diseases through the inhibition of fungal mycelial growth.

**DO NOT** apply using fogging or ultra low volume equipment.

**DO NOT** apply using handheld mistblowers or handheld airblast equipment.

MEDALLION Fungicide provides control of turf diseases when used in an integrated disease management program.

For optimum turfgrass quality and disease control, use MEDALLION Fungicide in conjunction with turfgrass management practices that promote plant health. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease is essential to determine the best control measures.

## **10.0 DIRECTIONS FOR USE**

### **10.1 General Information**

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**DO NOT** allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds, or other waters.

**DO NOT** apply more than 180 mL product/100 m<sup>2</sup>/year (18.0 L product/ha/year) on turfgrass.

## 10.2 Application Instructions

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

**DO NOT** apply using aerial application equipment.

## 10.3 Mixing Instructions

To prepare spray solution, fill half of the required water volume in the spray tank with clean water and begin agitation. Add the required amount of MEDALLION Fungicide to the tank, allowing time for good dispersion. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation. Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers should be calibrated before application and thoroughly cleaned immediately after application. Do not use silicone based products with MEDALLION Fungicide due to possible phytotoxicity.

## 10.4 MEDALLION Fungicide + Tank Mixtures

MEDALLION Fungicide is usually compatible with many tank-mix partners registered for use on turf.

If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used. Do not combine MEDALLION Fungicide in the spray tank with pesticides, surfactants, or fertilizers; unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use. To determine the physical compatibility of MEDALLION Fungicide with other products, use a jar test.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA (1-877-964-3682) for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

## 10.5 Buffer Zones

Use of the following spray methods or equipment **DOES NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats

(such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Turf		4	1	1	1	1
	Outdoor ornamentals		1	1	0	0	0
Airblast	Outdoor ornamentals	Early growth stage	20	2	0	0	0
		Late growth stage	10	1	0	0	0

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

## 11.0 CROP USE DIRECTIONS

The following use instructions will provide control of the listed diseases unless stated otherwise.

### 11.1 Turf: Golf Courses

TURF: GOLF COURSES	
DISEASES	<b>Anthracnose</b> ( <i>Colletotrichum cereale</i> ) <b>Brown patch</b> ( <i>Rhizoctonia solani</i> ) <b>Leaf spot</b> ( <i>Drechslera poae</i> )
PRODUCT RATE	6 L/ha or 60 mL/100 m <sup>2</sup>
APPLICATION INTERVAL	14 days
DISEASES	<b>Fusarium patch</b> ( <i>Microdochium nivale</i> )
PRODUCT RATE	3-6 L/ha or 30-60 mL/100 m <sup>2</sup>
APPLICATION DIRECTIONS	Use low rate on fairways only under conditions of low to moderate disease pressure.  For control of <i>Microdochium nivale</i> over winter (pink snow mould) high rate of 60 mL/100 m <sup>2</sup> rate is required. Refer to Pink snow mould ( <i>Microdochium nivale</i> ) section below for more information.
APPLICATION INTERVAL	14 days

<b>DISEASES</b>	<b>Brown ring patch (<i>Waitea circinata</i> var. <i>circinata</i>)</b>
<b>PRODUCT RATE</b>	3-6 L/ha or 30-60 mL/100 m <sup>2</sup>
<b>APPLICATION INTERVAL</b>	7-14
<b>MAXIMUM APPLICATIONS OF PRODUCT PER YEAR</b>	3
<b>APPLICATION DIRECTIONS</b>	<p>For best control, use as a component of an integrated disease management program.</p> <p>Begin applications when conditions are favourable for disease infection and prior to disease symptom expression.</p> <p>Under conditions of high disease pressure, use the highest label rate and the shortest interval. For optimum disease control tank mix or alternate MEDALLION Fungicide with a contact fungicide such as DICONIL® 2787 or DICONIL Ultrex.</p> <p>Apply in sufficient water to ensure thorough coverage. 4-8 litres of water per 100 m<sup>2</sup> is recommended.</p> <p>Use AI, XC or XR nozzles.</p> <p>Apply after mowing.</p> <p>Allow sprayed area to completely dry before irrigation.</p>
<b>DISEASE</b>	<b>Pink snow mould (<i>Microdochium nivale</i>)</b>
<b>PRODUCT RATE</b>	6 L/ha or 60 mL/100 m <sup>2</sup>
<b>MAXIMUM APPLICATIONS OF PRODUCT PER YEAR</b>	1
<b>APPLICATION INSTRUCTIONS</b>	<p>For optimum disease control tank mix MEDALLION Fungicide with A15457TO Fungicide. Under conditions of high disease pressure also include BANNER MAXX or Heritage MAXX in the tank mix.</p> <p>Make one application in the late fall before snow cover.</p> <p><b>DO NOT</b> apply on top of snow or when the temperature is below 0°C.</p> <p>For best control, use as a component of an integrated disease management program.</p> <p>Apply in sufficient water to ensure thorough coverage. 2-8 litres of water per 100 m<sup>2</sup> is recommended.</p> <p>Use AI, XR or Syngenta XC nozzles.</p> <p>Apply after mowing and do not mow after application.</p>

## 12.0 MINOR USES

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Syngenta Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

### DIRECTIONS FOR USE:

The following use instructions will provide control of the listed diseases unless stated otherwise.

#### 12.1 Ornamentals (Outdoor and Greenhouse)

ORNAMENTALS (Outdoor and Greenhouse)	
<b>DRENCH APPLICATION</b>	
<b>DISEASES</b>	Rhizoctonia stem rot and aerial blight ( <i>Rhizoctonia solani</i> ) Fusarium wilt ( <i>Fusarium oxysporum</i> ) - suppression only Black root rot ( <i>Thielaviopsis basicola</i> )
<b>PRODUCT RATE</b>	300 – 600 mL/1000 L of water
<b>APPLICATION METHOD</b>	Drench
<b>APPLICATION INTERVAL</b>	Completely drench the growing medium at planting then if needed apply 21 to 28 days later
<b>MAXIMUM APPLICATIONS OF PRODUCT PER GROWING CYCLE</b>	2
<b>APPLICATION DIRECTIONS</b>	<p>2 applications per growing cycle during conditions favourable for disease development are usually adequate to control diseases of ornamentals.</p> <p>Precaution for irrigation applications: Where distribution patterns do not overlap sufficiently, unacceptable control may result. Where distribution patterns overlap excessively, injury to desirable plants may result.</p> <p>Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.</p> <p>Plant safety has been found to be acceptable for many ornamental plants, however, not all possible plant species and varieties have been tested under all conditions. It is recommended to test MEDALLION Fungicide on a small portion of the crop to ensure that a phytotoxic response will not occur.</p> <p>DO NOT apply more than 3000 L/ha for drench application to greenhouse ornamentals.</p>

**ORNAMENTALS (Outdoor and Greenhouse)****FOLIAR APPLICATION**

<b>DISEASES</b>	<b>Alternaria leaf spot (<i>Alternaria alternata</i>, <i>A. zinnae</i>) Rhizoctonia stem rot and aerial blight (<i>Rhizoctonia solani</i>)</b>
<b>PRODUCT RATE</b>	300 – 600 mL/1000 L of water
<b>DISEASES</b>	<b>Botrytis blight/grey mould (<i>Botrytis cinerea</i>)</b>
<b>PRODUCT RATE</b>	600 – 1200 mL/1000 L of water
<b>DISEASES</b>	<b>Boxwood blight (<i>Cylindrocladium buxicola</i>)</b>
<b>PRODUCT RATE</b>	1200 mL/1000 L of water
<b>APPLICATION METHOD</b>	Foliar
<b>APPLICATION INTERVAL</b>	7-14 days
<b>MAXIMUM APPLICATIONS OF PRODUCT PER GROWING CYCLE</b>	4
<b>APPLICATION DIRECTIONS</b>	<p><b>DO NOT</b> apply more than four consecutive applications per growing cycle; rotate to another registered product with a different mode of action.</p> <p>Precaution for irrigation applications: Where distribution patterns do not overlap sufficiently, unacceptable control may result. Where distribution patterns overlap excessively, injury to desirable plants may result.</p> <p>Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.</p> <p>Plant safety has been found to be acceptable for many ornamental plants, however, not all possible plant species and varieties have been tested under all conditions. It is recommended to test MEDALLION Fungicide on a small portion of the crop to ensure that a phytotoxic response will not occur.</p> <p><b>DO NOT</b> apply more than one application at a spray volume of 1500 L/ha for foliar application to greenhouse cut flowers. A maximum of four applications can be made to greenhouse cut flowers if a reduced spray volume of 500 L/ha is used.</p>

## 12.2 Greenhouse Vegetables

<b>GREENHOUSE VEGETABLES</b>	
<b>CROP</b>	<b>Greenhouse Cucumber</b>
<b>DISEASES</b>	Suppression fusarium stem and root rot caused by <i>Fusarium oxysporum</i> f.sp. <i>radicis-cucumerinum</i>
<b>APPLICATION METHOD</b>	Drench application
<b>RATE OF APPLICATON</b>	250 mL solution per plant (solution: 300 mL product in 1000 L of water) (Equivalent to 0.75 L product/ha (93 g ai/ha) based on planting density for GH cucumber of 10,000 plants/ha)
<b>APPLICATION VOLUME</b>	Apply 250 mL of the Medallion mixture to the growing media at the base of each plant
<b>MAXIMUM APPLICATIONS OF PRODUCT PER GROWING CYCLE</b>	Apply 1 application per crop cycle
<b>APPLICATION TIMING</b>	Growing medium drench: apply to seedling crop prior to transplanting or transfer to larger containers. Completely drench growing medium with mixture. It is recommended to test MEDALLION alone and with any mixtures on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.
<b>PREHARVEST INTERVAL (PHI)</b>	1 day
<b>APPLICATION DIRECTIONS</b>	<p>Completely drench growing medium with mixture</p> <p><b>DO NOT</b> apply MEDALLION Fungicide to the foliage.</p> <p><b>DO NOT</b> use in the propagation house.</p> <p><b>DO NOT</b> apply to cucumbers intended for transplant out to fields.</p> <p>It is recommended to test MEDALLION alone and with any mixtures on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.</p>

<b>GREENHOUSE VEGETABLES</b>	
<b>CROP</b>	<b>Greenhouse Pepper</b>
<b>DISEASES</b>	Suppression of fusarium crown rot, stem rot and root rot caused by <i>Fusarium solani</i>
<b>APPLICATION METHOD</b>	Drench application
<b>RATE OF APPLICATON</b>	250 mL solution per plant (solution: 300 mL product in 1000 L of water) (Equivalent to 2.25 L product/ha (281 g ai/ha) based on planting density for GH pepper of 30,000 plants/ha)
<b>APPLICATION VOLUME</b>	Apply 250 mL of the Medallion mixture to the growing media at the base of each plant
<b>MAXIMUM APPLICATIONS OF PRODUCT PER GROWING CYCLE</b>	Apply 2 application per crop cycle.
<b>APPLICATION TIMING (CROP / PEST STAGE)</b>	Growing medium drench: apply to seedling crop prior to transplanting or transfer to larger containers. Completely drench growing medium with mixture. It is recommended to test MEDALLION alone and with any mixtures on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.
<b>PREHARVEST INTERVAL (PHI)</b>	1 day
<b>APPLICATION DIRECTIONS</b>	<p><b>DO NOT</b> apply Medallion Fungicide to the foliage.</p> <p><b>DO NOT</b> use in the propagation house.</p> <p><b>DO NOT</b> apply to peppers intended for transplant out to fields.</p> <p>Completely drench growing medium with mixture</p> <p>It is recommended to test MEDALLION alone and with any mixtures on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.</p>

<b>GREENHOUSE VEGETABLES</b>	
<b>CROP</b>	<b>Greenhouse Tomato</b>
<b>DISEASES</b>	<b>Suppression of fusarium wilt, incl. yellow stem, crown and root rot (<i>Fusarium oxysporum f. sp. radicis-lycopersici</i>, and <i>F. solani</i>)</b>
<b>APPLICATION METHOD</b>	Drench application
<b>RATE OF APPLICATION</b>	250 mL solution per plant (solution: 300 – 600 mL product/1000 L of water) (Equivalent to 15 – 45 mL product/100m <sup>2</sup> or 1.5 – 4.5 L product/ha (188 – 563 g ai/ha) based on planting density for GH tomato of 20,000 – 30,000 plants/ha)
<b>APPLICATION VOLUME</b>	Apply 250 mL of the MEDALLION mixture to the growing media at the base of each plant
<b>MAXIMUM APPLICATIONS OF PRODUCT PER GROWING CYCLE</b>	Apply 2 applications per crop cycle.
<b>APPLICATION INTERVAL</b>	Completely drench the growing medium at planting then if needed apply 21 to 28 days later
<b>APPLICATION TIMING (CROP / PEST STAGE)</b>	Growing medium drench; apply at transplant or thereafter prior to or at onset of disease when conditions are favourable in the production greenhouse
<b>PREHARVEST INTERVAL (PHI)</b>	1 day
<b>APPLICATION DIRECTIONS</b>	<p>2 applications per growing cycle during conditions favourable for disease development.</p> <p><b>DO NOT</b> apply MEDALLION Fungicide to the foliage.</p> <p><b>DO NOT</b> use in the propagation house.</p> <p><b>DO NOT</b> apply to tomatoes intended for transplant out to fields.</p> <p>Precaution for irrigation applications: Where distribution patterns do not overlap sufficiently, unacceptable disease management may result. Where distribution patterns overlap excessively, injury to desirable plants may result</p> <p>Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.</p> <p><b>DO NOT</b> enter or allow worker entry into treated areas during restricted entry interval (REI) of 12 hours.</p> <p>It is recommended to test MEDALLION alone and with any mixtures on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application.</p>

### 13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, MEDALLION Fungicide contains a Group 12 Fungicide. Any fungal

population may contain individuals naturally resistant to MEDALLION Fungicide and other Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.

Where possible, make use of predictive disease models to effectively time fungicide applications.

**To delay fungicide resistance:**

Where possible, rotate the use of MEDALLION Fungicide or other Group 12 fungicide with different groups that control the same pathogens.

Do not apply more than the maximum number of applications listed in the label.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at [www.syngenta.ca](http://www.syngenta.ca).