

# Mecoprop-P

## HERBICIDE

Mecoprop-P is a liquid herbicide for use on grasses (including turf, golf greens and fairways) and cereals. It controls tough broadleaved annual and perennial weeds.

**GUARANTEE:** Mecoprop-P 150 g/L (d-isomers equivalent present as potassium salt).

**CROPS:** Barley, bentgrasses, bluegrasses, oats, red fescue, rye grasses and wheat.

**RATE:** 5.5-8.5 L/ha (60-90 ml/100m<sup>2</sup>) (2.2-3.4 L/acre).

**WATER VOLUME:** Use 300-400 L/ha at a pressure not to exceed 275 kPa (27-36 gallons/acre). On small areas, apply at the recommended rates in 10 L of water/100 m<sup>2</sup>. In cereals, apply in 200-300 L/ha (18-27 gallons/acre).

**TANK MIXES:** On fairways and lawns, MCPA Amine 500 (3 L/ha) or 2,4-D (6.6 L/ha).

**TIMING:** For best results, apply to young, actively growing weeds in the 2-4 leaf stage. Treat wheat, oats and barley from the time the crop is in the 3-leaf expanded stage to the early flag leaf (shot blade) stage. Application can be made to seedling or established grasses.

### WEEDS CONTROLLED

**Grasses** – 5.5 L/ha for seedlings, 8.5 L/ha on established plants.

#### Susceptible weeds

Common chickweed

Mouse-eared chickweed

Clover

Plantain

#### Moderately susceptible – 8.5 L/ha

Black medic

Creeping charlie

Buttercup

Dandelion (with 2,4-D)

#### Wheat, barley and oats

#### Seedling 5.5 L/ha, mature 7 L/ha

Black medick

Corn spurry

Canada thistle  
(top growth only)

Lamb's quarters

Chickweed

Plantain

Cleavers

Stitchwort

Clovers (volunteer)

Wild mustard



## QUICK SHEET

**ALWAYS READ THE LABEL BEFORE USE**

**CAUTION:** Avoid spraying in very hot weather or drought conditions. Do not spray bentgrass when temperatures are above 27°C, particularly if high rates are used. Do not spray on grain underseeded to legumes. Do not apply by air.

**RAINFAST:** Rainfall within 4 - 6 hours may reduce effectiveness. Do not water grasses for 24 hours after treatment.

**PREHARVEST INTERVAL:** Do not graze the treated crops or cut for hay. Do not cut grass 3-4 days before or after treatment.

**RE-ENTRY TIME:** Allow thorough drying before re-entering.

**PACKAGE SIZE:** 2 x 10 L.

**RESIDUAL CONTROL:** None.

**ADDITIONAL:** Apply in warm weather under good growing conditions. Recommended water volumes are essential for optimum weed control. May act more slowly than some weed killers and 3 weeks may elapse before effect on weeds is apparent.

**SURFACTANT NEEDED:** None.

**RE-CROPPING:** No restrictions the year after application.

**STORAGE:** Do not freeze. If stored for 1 year or longer, shake well before using.

**HERBICIDE GROUP:** 4 Herbicide.

## NOTES:

[illegible]

**www.uap.ca**

**West: 1-800-561-5444    Ontario & Maritimes: 1-800-265-5444    Quebec 1-800-361-9369**